


WILD

GARDEN

POLLINATION



Early Blossoming
Apple Tree Flowers
(Early Stage)



{ The Far Orchard }
Apple Tree }

Flowers through
April to June




Common blue bell

Haiesinoidia non'scripta

wild garden at the barn

WILD CHERRY
Prunus avium




Flowers April-May

collected from the Wild Garden
at The Barn, Banchory

White Clover

Flowers June to September

The flowers are mainly collected by outbreeding insects and honey bees.




'Sweet Cicely'
Myrrhis odorata

Very common in Scotland
and the north of England.
Rarely seen south of
Birmingham.



April - June

Sweet Cicely was used to sweeten
food and drink before sugar came
to the British Isles.




Daffodil
wild gardens the barn
bees only go to daffodils
when there is nothing else
to offer.



May
Flowering

Dandelion
(*Taraxacum*)

Collected 4/5/24
The Barn, Banchory



Japanese kerria
wild garden the barn
kerria japonica

Ficaria verna
'lesser celandine'

When Aslan returned in C.S. Lewis's *The Lion, The Witch and the Wardrobe*, and the woodland turns from winter to spring, the ground was covered in all directions with yellow celandine.




part of the buttercup family

habitat:
woodland, grassland
gardens

flowering season:
January - April

As one of the first flowers to appear after winter, they provide an important nectar source for queen bumblebees



MOORNING WIDOW

COLLECTED THE BARN WILD GARDEN
MAY THE FOURTH BE WITH YOU
LOCATED BY SOUND
BEES DEEP WITHIN

Winter Heath

Erica carnea

flowering season: January to April



native to western, central, and southern Europe, as well as parts of Morocco. It can be found anywhere from the lowlands to high alpine regions.

Due to its early flowering time, winter heath is an important plant for bees when they first emerge after winter. And bees, in turn, are necessary for pollinating winter-flowering heather.

NETTLES URTICA is a GENUS
OF Flowering PLANT
IN the FAMILY
URTICACEAE

MANY Species have stinging hairs
and may be called nettles or
stinging nettles

Can Find them January to Dec

Common plant


Found in Gardens, hedgerows, fields,
woodlands and many habitats.

Preference for damp, Fertile and
disturbed ground makes it a
good coloniser of places
enriched by human activities
such as agricultural development



ROSE HIP

OR ROSEHUR
also called Rose
haw or rose haw
is a accessory fruit



04/05/2024
We collected it from
the wild flower
garden at the
barn.

Typically Red to
but orange but range
from dark purple to Black
In some species.

Rose hips begin to form after pollination
of flowers in Spring or Early Summer
and ripen in late Summer through to
Autumn.




IRISH SAXIFRAGE
THE BARN WILD GARDEN
INSECT HOTEL



COMPLEX
MASS OF
MANY LEAF
BUNCHES

POLLINATED BY FLIES + BEES
MAGENTA TO ROSE BLOOMS

4.5.2024



Irish Saxifrage

saxifraga rosacea



